



## Worked

## Solutions


Q1. i)  $f(x) = 2x - 3$




ii)  $g(x) = 3x^2 - 5x + 2$




iii)  $h(x) = e^x$




iv)  $f(x) = \sin x$



v)  $g(x) = \ln x$



vi)  $h(x) = |2x - 3|$



Q2.

i)  $f(x) = x^3 - 2x^2 + 3x, -3 \leq x \leq 4$

$(-3, -54), (-2, -22), (-1, -6), (0, 0)$   
 $(1, 2), (2, 6), (3, 18), (4, 44)$

ii)  $g(x) = \ln(2x - 3), 3 \leq x \leq 6$

$(3, 1.0986), (4, 1.61), (5, 1.95)$   
 $(6, 2.2)$

iii)  $h(\theta) = 2\sin 3\theta, -180 \leq \theta \leq 180^\circ$

As its  $3\theta$ , we "step" in  $30^\circ$   
 as the period will be  $\frac{360}{3} = 120^\circ \div 4 = 30^\circ$

$(-180, 0), (-150, -2), (-120, 0)$   
 $(-90, 2), (-60, 0), (-30, -2)$   
 $(0, 0), (30, 2), (60, 0)$   
 $(90, -2), (120, 0), (150, 2)$   
 $(180, 0)$

Q3.  $P(x) = 5x + 4, g(x) = 3x - 2$

i)  $f \circ g(9) \Rightarrow g(9)$  first

$g(9) = 3(9) - 2 = 27 - 2 = 25$

$f(25) = 5(25) + 4 = \boxed{129}$

ii)  $g^2(x) = g \circ g(x) \Rightarrow g(x)$  first

$g(x) = 3x - 2$

$g(3x - 2) = 3(3x - 2) - 2$

$= 9x - 6 - 2$

$= \boxed{9x - 8}$

Q4.  $f(x) = x - \frac{3}{2}$   $g(x) = \frac{2x-5}{3}$

i)  $fg(4) \Rightarrow g(4)$  first  
 $g(4) = \frac{2(4)-5}{3} = \frac{3}{3} = 1$

$f(1) = 1 - \frac{3}{2} = \boxed{-\frac{1}{2}}$

ii)  $gf(x) \Rightarrow f(x)$  first  
 $f(x) = x - \frac{3}{2}$   
 $g(x - \frac{3}{2}) = \frac{2(x - \frac{3}{2}) - 5}{3}$

$= \frac{2x - 3 - 5}{3}$   
 $= \boxed{\frac{2x-8}{3}}$

iii)  $f^2(x) = ff(x) \Rightarrow f(x)$  first  
 $f(x) = x - \frac{3}{2}$

$f(x - \frac{3}{2}) = (x - \frac{3}{2}) - \frac{3}{2}$   
 $= \boxed{x - 3}$

Q5.  $f(x) = e^{2x-1}$   $g(x) = \sqrt{x}$

i)  $fg(9) \Rightarrow g(9)$  first  
 $g(9) = \sqrt{9} = 3$  as  $x > 0$   
 $f(9) = e^{2(9)-1}$   
 $= \boxed{e^{17}}$

ii)  $gf(x) \Rightarrow f(x)$  first  
 $f(x) = e^{2x-1}$   
 $g(e^{2x-1}) = \boxed{\sqrt{e^{2x-1}}}$

iii)  $f^{-1}g(x) \Rightarrow g(x)$  first  
 $g(x) = \sqrt{x}$

To find  $f^{-1}(x)$ :

$y = e^{2x-1}$   
 $\ln y = \ln e^{2x-1}$

$\ln y = 2x - 1$

$1 + \ln y = 2x$

$\frac{1 + \ln y}{2} = x$

$\Rightarrow y = \frac{1 + \ln x}{2}$  (Swap  $x$  &  $y$ )

$\Rightarrow f^{-1}(x) = \frac{1 + \ln x}{2}$

$f^{-1}(\sqrt{x}) = \boxed{\frac{1 + \ln \sqrt{x}}{2}}$

Q6.  $f(x) = 6x - 3$   $g(x) = 5x + q$

$fg(x) \Rightarrow g(x)$  first

$g(x) = 5x + q$

$f(5x + q) = 6(5x + q) - 3$   
 $= 30x + 6q - 3$

$gf(x) \Rightarrow f(x)$  first

$f(x) = 6x - 3$

$g(6x - 3) = 5(6x - 3) + q$   
 $= 30x - 15 + q$

$fg(x) = gf(x)$

$30x + 6q - 3 = 30x - 15 + q$

$5q = -12$

$q = \boxed{\frac{-12}{5}}$

Q7.

$$f(x) = 6x - 1$$

$$\text{L.H.S.: } f[f(x + f(x))] ?$$

$$x + f(x) = x + 6x - 1 \\ = 7x - 1$$

$$f(7x - 1) = 6(7x - 1) - 1 \\ = 42x - 6 - 1 \\ = 42x - 7$$

$$\text{R.H.S.: } p[f(x)]$$

$$f(x) = 6x - 1$$

$$p[f(x)] = p(6x - 1) \\ = 6xp - p$$

$$\text{LHS} = \text{RHS}$$

$$42x - 7 = 6xp - p$$

Comparing  $x$  terms:

$$\Rightarrow 42x = 6xp$$

$$\boxed{p = 7}$$

Q8

$$f(x) = 3x - 5 \quad g(x) = 2x + 4$$

$$fg(x) \Rightarrow g(x) \text{ first}$$

$$g(x) = 2x + 4$$

$$f(2x + 4) = 3(2x + 4) - 5 \\ = 6x + 12 - 5 \\ = 6x + 7$$

$$fg(x) = 24$$

$$\Rightarrow 6x + 7 = 24$$

$$6x = 17$$

$$x = \boxed{17/6}$$

$$\text{Q9. } f(x) = 3^x \quad g(x) = x + 2$$

$$fg(x) \Rightarrow g(x) \text{ first}$$

$$g(x) = x + 2$$

$$f(x + 2) = 3^{x+2}$$

$$gf(x) \Rightarrow f(x) \text{ first}$$

$$f(x) = 3^x$$

$$g(3^x) = (3^x) + 2 = 3^x + 2$$

$$fg(x) = gf(x)$$

$$\Rightarrow 3^{x+2} = 3^x + 2$$

$$3^x \cdot 3^2 = 3^x + 2$$

$$9 \cdot 3^x = 3^x + 2$$

$$9 \cdot 3^x - 1 \cdot 3^x = 2$$

$$8 \cdot 3^x = 2$$

$$3^x = \frac{2}{8} = \frac{1}{4}$$

$$\log 3^x = \log \frac{1}{4}$$

$$x \cdot \log 3 = \log \frac{1}{4}$$

$$\Rightarrow x = \frac{\log \frac{1}{4}}{\log 3} = \boxed{-1.262}$$

Q10.

$$\text{i) } f(x) = x^2 + 8x - 3 \\ = \boxed{(x + 4)^2 - 19}$$

$$\text{ii) } g(x) = x^2 - 10x + 8 \\ = \boxed{(x - 5)^2 - 17}$$


$$\text{iii) } h(x) = x^2 + 12x + 43 \\ = \boxed{(x + 6)^2 + 7}$$

Q11. i)  $f(x) = 9x^2 + 36x - 4$   
 $= 9\left[x^2 + 4x - \frac{4}{9}\right]$   
 $= 9\left[(x+2)^2 - \frac{40}{9}\right]$   
 $= \boxed{9(x+2)^2 - 40}$

ii)  $g(x) = 4x^2 - 20x + 31$   
 $= 4\left[x^2 - 5x + \frac{31}{4}\right]$   
 $= 4\left[\left(x - \frac{5}{2}\right)^2 + \frac{3}{2}\right]$   
 $= \boxed{4\left(x - \frac{5}{2}\right)^2 + 6}$

Q12. i)  $f(x) = x^2 - 6x - 5$   
 $= \boxed{(x-3)^2 - 14}$

ii) Min pt =  $\boxed{(3, -14)}$   
 Axis of sym:  $\boxed{x=3}$



iv)  $y = (x-3)^2 - 14$   
 $y + 14 = (x-3)^2$   
 $x-3 = \sqrt{y+14}$   
 $x = \sqrt{y+14} + 3$   
 Swap  $x$  and  $y$ :  
 $y = \sqrt{x+14} + 3$

$\Rightarrow f^{-1}(x) = \boxed{\sqrt{x+14} + 3}$

Q13. i)  $f(x) = x^2 - 6x + 13$   
 $= \boxed{(x-3)^2 + 4}$

ii)  $y = (x-3)^2 + 4$   
 $y-4 = (x-3)^2$   
 $x-3 = \sqrt{y-4}$   
 $x = \sqrt{y-4} + 3$

Swap  $x$  and  $y$ :  
 $y = \sqrt{x-3} + 3$

$\Rightarrow f^{-1}(x) = \boxed{\sqrt{x-3} + 3}$

$\sqrt{\quad}$  can't be negative  
 $\Rightarrow x-3 \geq 0$

$\boxed{x \geq 3} = \text{Domain}$

iii) P o h  $\Rightarrow$  h first  
 $h(x) = x-3$


$P(x-3) = f(x)$   
 $P(x-3) = (x-3)^2 + 4$

$x-3 \rightarrow \boxed{\quad} (x-3)^2 + 4$

$\Rightarrow$  function must square the input and add 4


$\Rightarrow p(x) = \boxed{x^2 + 4}$

Q14.  $f(x) = 5x - 3 \quad \mathbb{R} \rightarrow \mathbb{R}$




- ii) Injective? Yes as any horizontal line in graph above will cut at most once  
 Surjective? Yes as any horizontal line in graph above will cut at least once.  
 Bijective? Yes as it is both injective and surjective

Q15.  $g(x) = x^2 - 2x \quad \mathbb{R} \rightarrow \mathbb{R}$



- ii) Injective? No as line  $y = 2$  cuts more than once.  
 Surjective? No as line  $y = -20$  never cuts the graph.  
 Bijective? No as it's neither injective nor surjective.


Q16.



- Injective? No as line  $y = \frac{1}{2}$  cuts twice in diag above.  
 Surjective? No as line  $y = -1.5$  never cuts the graph.  
 Bijective? No as it's neither injective nor surjective.

Q17.

$x$	-3	-2	-1	0	1	2	3
$x^2$	9	4	1	0	1	4	9



- ii) Not injective as line  $y = 3$  crosses twice.  
 As codomain is positive real numbers, it is surjective as all horizontal lines above the x-axis will cut the graph at least once.

iii) Domain changed to  $x \geq 0$  or  $x \leq 0$

Q18. i)  $f(x) = 5x - 2$

$$y = 5x - 2$$

$$5x = y + 2$$

$$x = \frac{y+2}{5}$$

Swap  $x$  and  $y$ :

$$y = \boxed{\frac{x+2}{5}}$$

ii)  $g(x) = x^2 - 6x$

$$y = x^2 - 6x$$

$$y = (x-3)^2$$

$$\sqrt{y} = x - 3$$

$$x = 3 + \sqrt{y}$$

Swap  $x$  and  $y$ :

$$y = \boxed{3 + \sqrt{x}}$$

iii)  $h(x) = 2(3^x)$

$$y = 2 \cdot (3^x)$$

$$\frac{y}{2} = 3^x$$

$$\log \frac{y}{2} = \log 3^x$$

$$x \cdot \log 3 = \log \frac{y}{2}$$

$$x = \frac{\log \frac{y}{2}}{\log 3}$$

Swap  $x$  and  $y$ :

$$y = \boxed{\frac{\log \frac{x}{2}}{\log 3}}$$

iv)  $P(x) = \sqrt{2x-4}$

$$(y)^2 = (\sqrt{2x-4})^2$$

$$2x - 4 = y^2$$

$$2x = y^2 + 4$$

$$x = \frac{y^2+4}{2}$$

Swap  $x$  and  $y$ :


$$y = \boxed{\frac{x^2+4}{2}}$$

v) Domain of  $p(x)$  was  $x \geq 2$  as  $\sqrt{\quad}$  can't be negative.

Range of  $p(x)$  was  $y \geq 0$  as you can't get a negative number out of a square root

$\Rightarrow$  Domain and range of  $p^{-1}(x)$  will be the opposite

$$\Rightarrow \boxed{\text{Domain: } x \geq 0 \text{ and Range: } y \geq 2}$$



Q19. i)  $f(x) = \frac{x-6}{4}$

$$y = \frac{x-6}{4}$$

$$4y = x - 6$$

$$x = 4y + 6$$

Swap  $x$  and  $y$ :

$$y = \boxed{4x + 6}$$

ii)  $g(x) = 4x + \frac{2}{3}$

$$y = 4x + \frac{2}{3}$$

$$y - \frac{2}{3} = 4x$$

$$x = \frac{y - \frac{2}{3}}{4}$$

$$= \frac{3y - 2}{12}$$

$$x = \frac{3y - 2}{12}$$

Swap  $x$  and  $y$ :

$$y = \boxed{\frac{3x - 2}{12}} \text{ or } \boxed{\frac{x - \frac{2}{3}}{4}}$$

iii)  $h(x) = \sqrt{x+4}$

$$(y)^2 = (\sqrt{x+4})^2$$

$$y^2 = x + 4$$

$$x = y^2 - 4$$

Swap  $x$  and  $y$ :

$$y = \boxed{x^2 - 4}$$

Q20.

i)  $(y)^2 = (\sqrt{x-3})^2$

$$y^2 = x - 3$$

$$x = y^2 + 3$$

Swap  $x$  and  $y$ :

$$y = \boxed{x^2 + 3}$$

Domain of  $f(x) = x \geq 3$

$\Rightarrow$  Range of  $f^{-1}(x) = \boxed{x \geq 3}$

ii)  $y = x^3$

$$x = \sqrt[3]{y}$$


Swap  $x$  and  $y$ :

$$y = \boxed{\sqrt[3]{x}}$$

Domain of  $g(x) = x \in \mathbb{R}$

$\Rightarrow$  Range of  $g^{-1}(x) = \boxed{x \in \mathbb{R}}$

Q21



Q22

